

PAPER



CrossMark
click for updates

Cite this: *RSC Adv.*, 2015, 5, 78941

Design of chiral urea-quaternary ammonium salt hybrid catalysts for asymmetric reactions of glycine Schiff bases†

Maximilian Tiffner,^a Johanna Novacek,^a Alfonso Busillo,^b Katharina Gratzler,^a Antonio Massa^{*b} and Mario Waser^{*a}

Bifunctional chiral urea-containing quaternary ammonium salts can be straightforwardly synthesised in good yield and with high structural diversity *via* a scalable and operationally simple highly telescoped sequence starting from *trans*-1,2-cyclohexanediamine. These novel hybrid catalysts were systematically investigated for their potential to control glycine Schiff bases in asymmetric addition reactions. It was found that Michael addition reactions and the herein presented aldol-initiated cascade reaction can be carried out to provide enantiomeric ratios up to 95 : 5 and good yields under mild conditions at room temperature.

Received 21st July 2015
Accepted 28th August 2015

DOI: 10.1039/c5ra14466c


www.rsc.org/advances

Introduction

Chiral onium salt (phase-transfer) catalysis is one of the fundamental non-covalent activation strategies in asymmetric catalysis.¹ Besides monofunctional chiral ammonium salt catalysts, the use of bifunctional derivatives has emerged as a powerful strategy for numerous applications.² Whereas the vast majority of such catalysts contain an additional OH-group as the second coordination site,^{2–4} the design and application of ammonium salts containing alternative H-bonding motifs has so far been less exhaustively investigated.^{2,5–7} While the groups of Lassaletta and Fernández,^{5a} Dixon,^{5b} Smith,^{5c} and Lin and Duan^{5d} introduced powerful Cinchona alkaloid-based (thio)-urea containing bifunctional ammonium salt catalysts, Zhao *et al.*⁶ recently developed a modular approach to access α -amino acid-based (thio)-urea/ammonium salt hybrid catalysts. Simultaneously, our groups introduced *trans*-1,2-cyclohexanediamine-based bifunctional ammonium salts **1** (Scheme 1).⁷ An extensive screening of different salts allowed us to identify **1a** as the most efficient chiral catalyst for asymmetric α -fluorination of β -ketoesters **2**.^{7a} In addition, catalyst **1b** was found to be promising for a newly developed aldol-initiated cascade reaction of glycine Schiff base **5** with cyanobenzaldehyde **6**.^{7b} In the initial optimisation of this powerful transformation only one cyclohexanediamine-based catalyst **1** was tested and it was also found that the Cinchona alkaloid-based catalyst **9** introduced by

Dixon^{5b} and Smith^{5c} gave the highest e.r. of 85 : 15 among the existing bifunctional ammonium salt motifs tested so far (Scheme 1, lower reaction).^{7b}

These encouraging initial results with this novel family of asymmetric catalysts prompted our groups to initiate a joint project focusing on the further development and exploration of



Scheme 1 Hybrid ammonium salts **1** and their applications reported previously.

^aInstitute of Organic Chemistry, Johannes Kepler University Linz, Altenbergerstraße 69, 4040 Linz, Austria. E-mail: mario.waser@jku.at; Tel: +43 732 2468 8748

^bDepartment of Chemistry and Biology, University of Salerno, 84084-Fisciano, Italy. E-mail: amassa@unisa.it; Tel: +39 089 969565


† Electronic supplementary information (ESI) available: Experimental details and copies of NMR spectra and HPLC analyses. See DOI: 10.1039/c5ra14466c

the hybrid salts **1** for stereoselective transformations. Thus we especially focused on optimisation of the existing synthesis route^{7a} to fuel the demand for larger quantities of the readily tuneable catalysts **1** and on systematic testing of these catalysts for asymmetric reactions of glycine Schiff bases **5** (*i.e.* the already mentioned cascade reaction and Michael addition reactions). Hereby a detailed screening of these catalysts to increase the selectivity in the cascade synthesis of compound **8** was considered to be particularly interesting, as our modular catalyst synthesis approach should allow us to overcome the limitations of the existing catalyst motifs. Interestingly, the groups of Liu and Soloshonok recently achieved a highly asymmetric synthesis of analogous lactams in a complementary approach by employing an asymmetric auxiliary-based approach using chiral glycine Schiff base Ni(II) complexes.⁸ This report once again proves the high potential of this robust auxiliary approach for the synthesis of chiral α -amino acid derivatives.⁹

Results and discussion

Catalyst synthesis

In contrast to recent reports by others who described incorporation of the H-bonding donor first, followed by a final quaternarization step,^{5,6} we have chosen to use an opposite assembly strategy, carrying out the quaternarization of mono-protected diamine **11** first, followed by deprotection and coupling with an iso(thio)cyanate (Scheme 2).^{7a} This strategy allowed us to overcome some of the challenges of the commonly used protocols,^{5,6} as side reactions of the nucleophilic heteroatoms of the H-bonding donor with the alkylating agent can be avoided, thus resulting in a broader and more functional group tolerant synthesis route.



Initial Protocol [Ref 7a]:
Two tedious purifications by column chromatography and limitations in the quaternarization!
(10–30% overall yield from **11**)

Optimized Protocol(s):
Final step column chromatography only, high yield and control of impurities in the quaternarization!
(40–50% yield from **11** on up to 3 mmol scale)

Scheme 2 Catalyst synthesis: (A) R^1CHO (1 eq.), $NaBH_4$ (1.5 eq.), MeOH/THF, r.t., 15 h (no purification, >90% conversion); (B) MeI (6 eq.), K_2CO_3 (1.1 eq.), DMF, 60 °C, 30 h (no purification, around 70% crude yield over 2 steps); (C) CH_2O (2 eq.), $Na(OAc)_3BH$ (2 eq.), DCE, r.t., 15 h (no purification, >90% conversion); (D) R^1CH_2Y , DMF, 60 °C, 30 h (no purification, 60–80% crude yield over 2 steps); (E) HI (10 eq.), DCM, r.t., 2 h (extractive work-up with Na_2CO_3); (F) R^2NCX (1.2 eq.), DCM, r.t., 2–6 h (simple purification by column chromatography, 60–80% yield over 2 steps). X = O or S and Y = I, Br or Cl.

As outlined in Scheme 2, the catalysts **1** can be obtained from the known Boc-protected diamine **11**¹⁰ in four chemical steps. First, the introduction of bulkier (aromatic) residues for R^1 on the ammonium side can be easily accomplished by means of a reductive amination, giving the corresponding secondary amines **12** in high yields (>90% conversion) and without the need for any further purification. However, the subsequent quaternarization (step B, Scheme 2) was found to be the major bottleneck in the initial protocol.^{7a} It was not possible to introduce any residues that are sterically more demanding than a methyl group, thus allowing the syntheses of dimethyl-containing ammonium salts **13** only. In addition, we found that this exhaustive methylation is strongly dependent on the nature of substituent R^1 . While smaller, electron rich or electron neutral groups like a phenyl group allowed us to obtain **13** in slightly more than 50% yield starting from **11**, electron-poor or sterically more demanding groups performed significantly worse (down to around 25% yield). Hereby we faced two problems: first the final quaternarization was rather slow, leading to mixtures of *tert*-amine intermediates and the quaternary ammonium salts even after long reaction times. In addition, under the rather forcing and prolonged conditions we observed formation of large quantities of the trimethyl ammonium salt **13b** ($R^1 = H$). Formation of the latter can be explained by a nucleophilic substitution of the benzylic ammonium groups of compound **13** (*e.g.* by the base or the iodide counter anion) under the harsh reaction conditions, giving the more reactive dimethyl-containing *tert*-amine, which was then further methylated. This effect lowered the overall yields significantly and also made purification of the ammonium salts rather difficult. In addition, when using sterically even more demanding R^1 residues like an anthracenyl group no product **13** could be obtained.

To overcome these significant limitations, we first tested alternative (sterically less demanding) *N*-protecting groups (*e.g.* alloc) but with no success. Also other methylation agents did not improve the outcome. Gratifyingly, after a very careful screening of different conditions we finally found that carrying out the reaction with MeI (6 eq.) and solid K_2CO_3 (1.1 eq.) in DMF (60 °C) gave the target ammonium salts (**13**) in reliably high yields (around 70% *in situ*) and sufficient purity to be telescoped further in the sequence without any purification. It is noteworthy that almost no *tert*-amine intermediates and no trimethyl ammonium salt **13b** were formed (traces of **13b** originating from methylation of remaining **11** could be separated in the next step). This strategy was found to be very robust for when R^1 is electron-rich, -neutral, and -poor, giving access to derivatives that were not accessible by the initial route (*e.g.* 3,5-(CF_3)– C_6H_3 –). Unfortunately, when using sterically demanding R^1 groups (*e.g.* naphthyl or *ortho*-substituted aromatics) this route was still not very satisfactory (hereby the reactions mainly stalled at the *tert*-amine intermediates). However, the ammonium salts could be obtained in reasonable yields by first carrying out the dimethylation of **11** (step C) and then a final quaternarization with the appropriate benzylic halide (step D). Again the products were obtained in sufficient purity for direct further use. Thus, these newly developed quaternarization

conditions allowed us to obtain a much more diverse assembly of ammonium salts **13** which were then transformed into the catalysts **1** in two more steps. The only limitations in the present synthesis are the fact that only one sterically demanding group R^1 can be introduced and that groups bigger than a naphthyl group (e.g. anthracenyl) can only very slowly be incorporated.

The Boc-deprotection (step E) was initially carried out with TFA and the resulting salt (obtained by evaporation of the TFA and the solvent) was directly used for the final coupling step. However, this deprotection procedure gave mixtures of ammonium trifluoroacetates and iodides, which showed different catalytic properties and were difficult to separate. Use of HI for the deprotection was found to be beneficial for this reaction and, accompanied with an extractive work up, yielded the free amines **14** in good purity and yield. The extraction also allowed us to remove residual trimethyl ammonium salt **14b** ($R^1 = H$), which is significantly more hydrophilic than the aryl-containing derivatives. With this optimized procedure in hand, the final coupling could be carried out straightforwardly with different iso(thio)cyanates, requiring only one final (and simple) purification by silica gel column chromatography. Thus, a diversified assembly of different catalysts **1** can now be obtained in a telescoped and operationally simple manner and with satisfying overall yields (>30% based on **10**) on a practical scale (up to 3 mmol).

Asymmetric reactions of the glycine Schiff bases

The first transformation that we carefully investigated was the Michael addition of glycine Schiff base **5** to acceptors like methyl acrylate **15a**. This has been a thoroughly investigated reaction in asymmetric non-covalent organocatalysis in the past and it is noteworthy that most of the reported catalysts have their own characteristics with respect to their application scope (especially when using β -substituted acceptors).^{11–13} Thus we were curious to see whether our hybrid catalysts can be used (and systematically optimised) to control this important transformation. Table 1 gives an overview of the most significant results obtained from a very detailed screening of different catalysts and reactions conditions. We first identified the combination of toluene and solid CS_2CO_3 (1.5 eq.) as the best-suited solvent–base system for this reaction. Larger amounts of base favoured the racemic background reaction and non-aromatic solvents or aqueous (alternative) bases generally gave significantly lower selectivities (these effects were also carefully double-checked once the most active catalyst was identified).


With the solvent and base conditions set, we focused on the identification of the most active catalyst. All the initial reactions were run for 24 h using the same setup (Schlenk flask, stirring rate, and dilution) and stoichiometric ratio of the reagents to assure reproducible and comparable biphasic reaction conditions.

It immediately became apparent that ureas are better-suited than thioureas (see entries 1 and 2) and that the use of benzylic ammonium salts is beneficial (entry 3 vs. 1). Thus a series of hybrid catalysts with different aryl groups (R^1) on the

ammonium side (keeping the phenyl-urea group R^2 unchanged) were tested under identical conditions (entries 3–8 give the most significant results). Contrary to our results obtained in the α -fluorination of ketoesters^{7a} (shown in Scheme 1), the introduction of bulky naphthyl groups as R^1 was found to have no beneficial effect. In contrast, increasing the bulk by incorporating a *t*-butyl group on the aryl moiety (entry 6) significantly reduced the reaction rate. Also, the introduction of nitro substituents did not allow us to increase the selectivity (entry 7). Luckily, as already discussed above, the new synthesis protocol (Scheme 2) allowed us to introduce trifluoromethyl-substituted aryl groups as R^1 (i.e. 3,5-(CF_3)₂- C_6H_3) in good yield. This catalyst modification turned out to be the most fruitful amongst all the tested R^1 groups, giving **16a** with reasonable selectivity (e.r. = 84 : 16) and in high yield under the standard conditions (entry 8). Based on this encouraging result, we next systematically modified the urea-substituent R^2 . The presence of electron-withdrawing groups led to reduced selectivities in the case of CF_3 , nitro-, or diester-containing R^2 groups (entries 9–11). It is noteworthy that the latter was found to be the urea-modification of choice in our recent α -fluorination protocol.^{7a} Accordingly, these results show once more that such reactions always require a detailed and systematic screening of differently modified catalysts, thus illustrating the need for a flexible and functional group-tolerant catalyst synthesis as outlined in Scheme 2.


Interestingly, at this stage we found that replacing the aryl moieties on the urea side by incorporating a cyclohexyl group as R^2 instead resulted in a good selectivity of 87 : 13 (entry 12). Based on the fact that aliphatic groups were never found to be promising for any of our previously investigated reactions (published^{7a} or unpublished) this result came as a big surprise. By testing the influence of the reaction temperature we found that the selectivity can be increased by lowering the temperature, however this is accompanied by a reduced reaction rate (entries 14 and 15). Again, the reduced selectivity when using thiourea-moieties was proven by testing the analogous cyclohexyl-thiourea catalyst (entry 16). Finally, additional modifications introducing different alkyl groups as R^2 allowed us to identify an ethyl group as the most powerful urea-substituent, giving **16a** with an enantiomeric ratio of 89 : 11 under the standard conditions at room temperature. Further fine-tuning of the reaction conditions showed that the selectivity increases with higher dilution, however this results in a stepwise decrease of conversion (entries 18–23). The highest selectivities were obtained either by carrying out the reaction at a 0.02 M concentration of **5** at 25 °C (entry 21) or at higher concentration (0.08 M) at 0 °C (entry 20), albeit with reduced conversion. Furthermore, lowering the catalyst loading resulted in a reduced catalytic performance (entry 22), and further dilution did not allow us to improve the selectivity any further (entry 23).

To prove the necessity of the bifunctional nature of the catalysts **1** we also tested the simplified catalysts **17** and **18**, which performed with much less selectivity or did not give any product at all under the optimized reaction conditions (Scheme 3).

Table 1 Identification of the most active catalyst **1** and the best-suited reaction conditions for the addition of **5** to **15a**


No	R ¹	R ²	X	Conc. ^a [M]	T [°C]	Yield ^b [%]	e.r. ^c (R : S)
1	H	Ph	O	0.17	25	60	67 : 33
2			S			69	60 : 40
3	Ph		O			90	76 : 24
4	α -Np					72	78 : 22
5	β -Np					35	74 : 26
6	4- <i>t</i> Bu-C ₆ H ₄					6	79 : 21
7	4-NO ₂ -C ₆ H ₄					40	74 : 26
8	3,5-(CF ₃) ₂ -C ₆ H ₃					85	84 : 16
9		3-NO ₂ -C ₆ H ₄				83	81 : 19
10		3,5-(CO ₂ Me) ₂ -C ₆ H ₃				85	75 : 25
11		3,5-(CF ₃) ₂ -C ₆ H ₃				95	75 : 25
12		Cy				88	87 : 13
13					40	90	81 : 19
14					0	70	92 : 8
15					-20	5	92 : 8
16			S		25	91	65 : 35
17		<i>t</i> Bu	O			94	85 : 15
18		Et				96	89 : 11
19				0.08		87	90 : 10
20					0	45	95 : 5
21	3,5-(CF ₃) ₂ -C ₆ H ₃	Et	O	0.02	25	85	95 : 5
22 ^d						74	88 : 12
23				0.01		8	95 : 5

^a Based on **5**. ^b Isolated yield. ^c Determined from HPLC using a chiral stationary phase.¹¹ ^d Using 5 mol% catalyst.




Scheme 3 Control experiments using the simplified catalysts **17** and **18**.

Having identified the best-suited catalyst and the optimum reaction conditions (see entry 21, Table 1) we next investigated the application scope for this reaction (Scheme 4). One important fact that should be mentioned is that this biphasic transformation was found to be rather sensitive (e.r. and conversion) with respect to changes in the reaction setup (*i.e.* shape of the Schlenk flasks used and stirring rate). Therefore, when investigating the application scope we always carried out each test

reaction on a 0.1 mmol scale (based on **5**) with exactly the same setup and for comparison also on a smaller scale (0.02 mmol). All the results shown in Scheme 4 were thus carefully double-checked and found to be reproducible in at least two runs with two different setups each.

Testing different esters of the glycine Schiff base **5** first, we found that all gave good conversion within 24 h, but that the selectivity depends on the nature of the ester group, with bulky *t*-butyl esters of **5** giving the highest enantioselectivity in the addition to methyl acrylate **15a** (see results for products **16a–16c**). Interestingly, by changing the ester group of the Michael acceptor **15** both the conversion rate¹⁴ and the enantioselectivity were influenced (see products **16d–16f**). This was most notable in the synthesis of the di-*t*-butyl containing diester **16e** (only 15% conversion under standard conditions and a reduced e.r. of 83 : 17). Here the conversion could be increased by using 5 equivalents of base without affecting the e.r. (when using other acceptors the use of a larger excess of base usually resulted in a reduced e.r. because of an increased rate of the racemic background reaction). When testing phenyl vinyl sulfone as an acceptor (giving product **16g**) the selectivity dropped significantly, whereas *N,N*-dimethyl acrylamide and methyl vinyl ketone could be employed with reasonable selectivities (**16h** and **16i**). Unfortunately, the acrylamide acceptor was found to



Scheme 4 Application scope for the Michael addition of the glycine Schiff bases **5** to different Michael acceptors (**15**) using bifunctional catalyst **1c**. The configuration of products **16a**, **16b**, **16d–16g**, **16i**, **16j'**, **16k** and **16m** was determined by comparison of the analytical details with literature data,^{11–13,16,17} whereas the other products (**16c**, **16h**, **16j** and **16l**) were assigned by analogy.

react rather slowly and in this case the use of more base did not really allow us to overcome this limitation as the selectivity was clearly affected (this substrate was also very difficult to control in an enantioselective manner in a recent project using structurally different TADDOL-based PTCs¹¹).


Interesting results were obtained using prochiral electrophiles **15j–15m** (as shown in the lower part of Scheme 4). Using dimethyl maleate **15j**, the product **16j** could be obtained in good yield and with high diastereo- and enantioselectivity. It is noteworthy that Lambert and co-workers recently found that this substrate was unreactive when using their otherwise very powerful and highly selective cyclopropenimine chiral base catalysts.¹² Using dimethyl fumarate **15j'** instead gave the diastereomeric **16j'** in good yield and diastereoselectivity albeit with a slightly lower enantioselectivity. Another striking difference to the chiral cyclopropenimine base catalysis was also observed when employing *s-trans* acceptors **15k** and **15l**. Although cyclohexenone **15k** underwent a slow 1,4-addition under the standard conditions (which can be explained by a competing dimerization under basic phase-transfer

conditions¹⁵), the stereoselectivity obtained is still reasonably high.¹⁶ Even more interestingly, when using cyclopentenone **15l** we were able to obtain product **16l** in high yield and with good stereoselectivity.¹⁷ Finally, chalcone was also accepted well in this reaction, providing the product **16k** with reasonable selectivity and in good yield. Here it is of course fair to mention that for this substrate again Lambert's chiral base catalyst was reported to be more selective¹² and therefore the results obtained hereby may illustrate that our catalysts can provide a complementary and useful activation platform for future asymmetric organocatalytic transformations especially for addition reactions to *s-trans* Michael acceptors.

Having shown that the catalysts **1** can be systematically fine-tuned to obtain high selectivities for the Michael addition of the Schiff bases **5** to different Michael acceptors **15**, we next addressed the recently developed aldol-initiated cascade reaction of **5** with cyanobenzaldehyde **6**.^{7b} As already discussed above, the initial screening of existing catalysts showed that the Cinchona alkaloid-based bifunctional ammonium salt **9** introduced by Dixon^{5b} and Smith^{5c} gave the highest selectivity with a d.r. = 4 : 1 and an e.r. of 85 : 15 for the major diastereomer (see Scheme 1). With our optimized synthesis for the catalysts **1** in hand we put our efforts into identifying an even more selective catalyst for this powerful transformation. Table 2 gives an overview of the most significant results obtained thereby. The screening was carried out under the previously developed conditions^{7b} using solid K_2CO_3 as the base in CH_2Cl_2 (other solvents and bases were tested too but found to be less useful). In addition, initial tests showed that again ureas are more active and selective than thioureas and thus the optimization was carried out with ureas only.

Entry 1 shows the initially reported result^{7b} using the trimethylammonium-based catalyst **1b** and it was soon found that variation of the urea group alone does not significantly change the catalytic performance (entries 1–4). Introducing electron-neutral or bulky aryl groups on the ammonium side (entries 5–8) did not result in any improvement either. In contrast, the introduction of a naphthyl group even reduced the diastereo- and enantioselectivity (entry 8). Also the presence of aliphatic groups on the urea side (which was beneficial for our Michael reaction) did not allow us to improve the catalyst performance (entry 6) and a similar selectivity was obtained upon incorporation of more electron-rich aryl groups as R^1 (entries 9 and 10).

In sharp contrast, the introduction of electron-withdrawing groups on the aryl group R^1 on the ammonium side accompanied by the presence of an electron-poor aryl moiety R^2 on the urea side resulted in significantly superior catalysts (entries 11–13). First, the dinitro-substituted catalyst used in entry 11 almost matched the result previously obtained with the Cinchona catalyst **9**, both in terms of reactivity and selectivity. Gratifyingly, as already observed for the Michael addition, the introduction of a di-trifluoromethyl-phenyl group as R^1 again significantly improved the yield and the selectivity in this transformation (entries 12 and 13). Modifying the R^2 group finally resulted in the most promising catalyst, shown in entry 12, which gave product **8** in 83% isolated yield and with a good

Table 2 Identification of the most active catalyst **1** for the cascade reaction of **5** and **6**

No	R ¹	R ²	t [h]	Yield ^a [%]	d.r. ^b	e.r. major (minor) ^c
1	H	3,5-(CF ₃) ₂ -C ₆ H ₃	24	82	1.5 : 1	77 : 23 (60 : 40)
2		3-NO ₂ -C ₆ H ₄	24	66	1.4 : 1	77 : 23 (60 : 40)
3		4-CF ₃ -C ₆ H ₄	48	63	1.3 : 1	76 : 24 (51 : 49)
4		Ph	48	52	1.2 : 1	80 : 20 (59 : 41)
5	Ph		48	52	1 : 1	77 : 23 (52 : 48)
6		Cy	72	68	2 : 1	67 : 33 (51 : 49)
7		4-CF ₃ -C ₆ H ₄	48	60	1 : 1	80 : 20 (57 : 43)
8	β-Np	Ph	48	52	1 : 1	77 : 23 (54 : 46)
9	3,5-F ₂ -4-MeO-C ₆ H ₂	Ph	96	13	1.1 : 1	70 : 30 (57 : 43)
10	4-MeO-C ₆ H ₃	3-NO ₂ -C ₆ H ₄	48	59	1.3 : 1	77 : 23 (59 : 41)
11	3-NO ₂ -C ₆ H ₄	3-NO ₂ -C ₆ H ₄	26	65	2.4 : 1	85 : 15 (63 : 37)
12	3,5-(CF ₃) ₂ -C ₆ H ₃	3-NO ₂ -C ₆ H ₄	24	83	4 : 1	95 : 5 (75 : 25)
13	3,5-(CF ₃) ₂ -C ₆ H ₃	3,5-(CF ₃) ₂ -C ₆ H ₃	24	79	3 : 1	92 : 8 (73 : 27)


^a Isolated yield of **8**. ^b Determined from the ¹H NMR of the crude product. ^c Determined from HPLC using a chiral stationary phase.^{7b}

d.r. of 4 : 1 and excellent enantioselectivity for the major diastereoisomer (e.r. = 95 : 5; the minor diastereoisomer was always obtained with a lower enantiomeric ratio and no further improvement was possible).

Determination of the relative and absolute configuration of **8** turned out to be rather difficult as both diastereomers only crystallized as amorphous solids and thus X-ray structural analysis was not possible (the same was the case for the other derivatives shown in Scheme 5). The coupling constants of **7**

and **8** also did not enable us to unambiguously determine the relative configuration (NOESY experiments were also not clear). However, it was possible to assign the absolute configuration of the amino acid stereogenic center of the major diastereomer to be (*R*) by analysis of the Mosher amide derivatives of **8a**.¹⁸ Subsequent NMR studies of the compounds then indicated that the major diastereomer is most probably the *anti* isomer as depicted in Table 2.¹⁹

With the most active catalyst **1d** in hand we quickly tested two halide- and one methoxy-containing cyanobenzaldehydes (**6**) under the same conditions and found that these substrates were also well tolerated, albeit the stereoselectivities slightly decreased in these cases, especially for the minor diastereoisomers (Scheme 5). Nevertheless, a first proof-of-principle for the generality of this reaction was clearly made, thus illustrating the potential of these catalysts for such cascade-reactions and providing a complementary approach to the recently reported analogous auxiliary-based protocol.⁸

Scheme 5 Use of the cyanobenzaldehydes **6** for the cascade reaction.

Experimental

General information

¹H and ¹³C-NMR spectra were recorded on a Bruker Avance III 300 MHz spectrometer, on a Bruker Avance III 700 MHz spectrometer with a TCI cryoprobe or on Bruker DRX 400, 300 or 250 MHz spectrometers. All NMR spectra were referenced to the solvent peak. High resolution mass spectra were obtained using

an Agilent 6520 Q-TOF mass spectrometer with an ESI source and an Agilent G1607A coaxial sprayer or using a Thermo Fisher Scientific LTQ Orbitrap XL with an Ion Max API Source. Additional mass spectral analyses were carried out using an electrospray spectrometer, a Waters 4 micro quadrupole. Elemental analyses were performed with FLASHEA 1112 series-Thermo Scientific for CHNS-O apparatus. IR spectra were recorded on a Bruker Tensor 27 FT-IR spectrometer with an ATR unit. HPLC analyses were performed either by using a Waters instrument or using a Dionex Summit HPLC system with a Chiralcel OD-H (250 × 4.6 mm, 5 μm), a Chiralcel OD-R (250 × 4.6 mm, 10 μm), or a Chiralpak AD-H (250 × 4.6 mm, 5 μm) chiral stationary phase. Optical rotations were recorded on a PerkinElmer Polarimeter Model 241 MC and on a Schmidt + Haensch Polarimeter Model UniPol L 1000. All chemicals were purchased from commercial suppliers and used without further purification unless otherwise stated. All reactions were performed under an Ar atmosphere.

A detailed experimental section including all the procedures and analytical data of the catalysts and asymmetric reaction products as well as copies of the NMR spectra and HPLC chromatograms can be found in the ESI.†

Synthesis of catalysts 1c and 1d

Step 1 (step A in Scheme 2). 3,5-bis(Trifluoromethyl)-benzaldehyde (484 mg, 2 mmol) was added to a solution of **11** (428 mg, 2 mmol) (prepared from (*S,S*)-cyclohexanediamine dihydrochloride in analogy to the literature⁴⁰) in 10 mL of THF-methanol (1 : 1) and the solution was stirred at r.t. overnight. After addition of NaBH₄ (114 mg, 3 mmol, 1.5 eq.), stirring was continued for another 2 h at r.t. The reaction was quenched by addition of water and extracted with water/diethyl ether. The organic phase was washed with brine, dried over Na₂SO₄, and evaporated to dryness to obtain the crude product **12c** in almost quantitative yield (95%, 824 mg, 1.9 mmol), which could be directly used without any purification. Compound **12c** (R¹ = 3,5-(CF₃)₂-C₆H₃): ¹H NMR (300 MHz, δ, CDCl₃, 298 K): 1.08–1.35 (m, 4H), 1.42 (s, 9H), 1.61–1.85 (m, 2H), 1.96 (s (b), 1H), 1.99–2.15 (m, 2H), 3.31–3.47 (m, 1H), 3.83 (d, 1H, *J* = 14.1 Hz), 4.00 (d, 1H, *J* = 14.1 Hz), 4.48 (d, 1H, *J* = 7.3 Hz), 7.73 (s, 1H), 7.82 (s, 2H) ppm.

Step 2 (B). The amine **12c** (1.7 mmol) was dissolved in 3 mL of DMF. After addition of K₂CO₃ (287 mg, 2.1 mmol, 1.2 eq.) and methyl iodide (646 μL, 10.4 mmol, 6 eq.), the suspension was stirred for 30 h at 60 °C. After removal of excess methyl iodide under reduced pressure, the suspension was extracted with dichloromethane/brine. The organic phase was dried over Na₂SO₄ and removal of the solvent under reduced pressure gave **13c** (70%, 1.13 g, 1.2 mmol), which was used without further purification. Compound **13c** (R¹ = 3,5-(CF₃)₂-C₆H₃): [α]_D²¹ (*c* = 1.8, CH₂Cl₂) = −7.4°; ¹H NMR (500 MHz, δ, CDCl₃, 298 K): 1.32–1.40 (m, 1H), 1.46 (s, 9H), 1.58–1.66 (m, 1H), 1.66–1.74 (m, 1H), 1.77–1.84 (m, 1H), 1.95–2.06 (m, 3H), 2.59–2.64 (m, 1H), 3.22 (s, 3H), 3.28 (s, 3H), 4.11–4.18 (m, 1H), 4.93–5.00 (m, 1H), 5.16 (d, 1H, *J* = 12.8 Hz), 5.36 (d, 1H, *J* = 12.8 Hz), 5.87 (d, 1H, *J* = 9.9 Hz), 8.01 (s, 1H), 8.13 (s, 2H) ppm; ¹³C NMR (126 MHz, δ, CDCl₃,

298 K): 24.8, 24.8, 27.7, 28.5, 35.7, 49.7, 51.1, 51.7, 63.1, 77.8, 81.5, 122.8 (q, *J* = 273 Hz), 124.9, 130.3, 133.0 (q, *J* = 34 Hz), 133.6, 155.9 ppm; ¹⁹F NMR (282 MHz, δ, CDCl₃, 298 K): −62.8 ppm; IR (film): $\bar{\nu}$ = 3431, 3270, 3011, 2980, 2939, 2867, 1695, 1625, 1516, 1467, 1455, 1393, 1370, 1323, 1281, 1242, 1176, 1138, 1048, 1024, 904, 870, 844, 737, 709, 683 cm^{−1}; HRMS (ESI) *m/z* calcd for C₂₂H₃₁F₆N₂O₂⁺: 469.2284 [M⁺], found: 469.2281.

Step 3 (E). The ammonium salt **13c** (1.1 mmol) was dissolved in 12 mL of dichloromethane and hydroiodic acid (57 wt% aq. solution, 1.45 mL, 11 mmol, 10 eq.) was added. After stirring for 2 h at r.t., the reaction was basified with saturated Na₂CO₃ solution and extracted with dichloromethane. The organic phase was dried over Na₂SO₄ and removal of the solvent under reduced pressure gave crude **14c** in quantitative yield (689 mg, 1.1 mmol). Compound **14c** (R¹ = 3,5-(CF₃)₂-C₆H₃): ¹H NMR (300 MHz, δ, CDCl₃, 298 K): 1.31–1.76 (m, 5H), 1.84–1.95 (m, 1H), 1.96–2.07 (m, 1H), 2.27–2.37 (m, 1H), 3.06 (s, 3H), 3.17 (s, 3H), 3.46 (s (b), 2H), 3.56–3.68 (m, 1H), 4.06–4.17 (m, 1H), 5.12 (d, 1H, *J* = 12.7 Hz), 5.48 (d, 1H, *J* = 12.7 Hz), 7.96 (s, 1H), 8.16 (s, 2H) ppm.

Step 4 (F). A solution of **14c** and the corresponding isocyanate (1.2 eq.) in dichloromethane (20 mL per mmol of **14c**) was stirred for 4 h at r.t. Evaporation of the solvent under reduced pressure gave crude **1**, which was further purified by column chromatography (dichloromethane : methanol = 50 : 1 to 10 : 1) to obtain pure catalyst **1** in the reported yields.

Catalyst 1c. Obtained in 65% yield (123 mg, 0.217 mmol starting from 0.33 mmol of **13c**) as a colourless oil. [α]_D²¹ (*c* = 1.3, CHCl₃) = 13.0°; ¹H NMR (300 MHz, δ, CDCl₃, 298 K): 1.16 (t, *J* = 7.2 Hz, 3H), 1.24–1.41 (m, 1H), 1.43–1.66 (m, 2H), 1.67–1.87 (m, 2H), 1.90–2.12 (m, 2H), 1.49–1.61 (m, 1H), 3.06 (s, 3H), 3.18–3.34 (m, 5H), 4.22–4.38 (m, 1H), 4.59–4.52 (m, 1H), 5.32–5.47 (m, 2H), 5.99 (s, 1H), 6.92 (d, *J* = 9.7 Hz, 1H), 7.97 (s, 1H), 8.00 (s, 2H); ¹³C NMR (75 MHz, δ, CDCl₃, 298 K): 15.5, 24.7, 25.1, 27.4, 35.2, 35.9, 48.1, 50.9, 51.0, 65.3, 78.0, 122.7 (q, *J* = 273 Hz), 124.8, 130.5, 133.0 (q, *J* = 34 Hz), 133.4, 157.7 ppm; ¹⁹F NMR (282 MHz, δ, CDCl₃, 298 K): −62.9 ppm; IR (film): $\bar{\nu}$ = 3295, 3021, 2988, 2936, 2864, 2349, 2288, 1656, 1546, 1449, 1373, 1278, 1174, 1130, 904, 843, 751, 719, 682, 663, 593, 463, cm^{−1}; HRMS (ESI): *m/z* calcd for C₂₀H₂₈F₆N₃O⁺: 440.2131 [M⁺], found: 440.2118.

Catalyst 1d. Obtained in 67% yield (96 mg, 0.14 mmol, starting from 0.22 mmol of **13c**) as an orange oil. [α]_D²¹ (*c* = 0.75, CH₂Cl₂) = −29.3°; ¹H NMR (300 MHz, δ, CDCl₃, 298 K): 1.29–1.46 (m, 1H), 1.56–2.06 (m, 5H), 2.11–2.23 (m, 1H), 2.55–2.67 (m, 1H), 3.19 (s, 3H), 3.28 (s, 3H), 4.40–4.62 (m, 2H), 5.37 (s, 2H), 7.39 (t, 1H, *J* = 8.2 Hz), 7.47 (d, 1H, *J* = 9.2 Hz), 7.73 (dd, 2H, *J*₁ = 8.1 Hz, *J*₂ = 1.5 Hz), 7.84 (dd, 1H, *J*₁ = 8.1 Hz, *J*₂ = 1.8 Hz), 7.94 (s, 1H), 8.03 (s, 2H), 8.69 (t, 1H, *J* = 2.1 Hz), 9.11 (s, 1H) ppm; ¹³C NMR (75 MHz, δ, CDCl₃, 298 K): 24.5, 25.0, 27.3, 36.0, 49.1, 50.6, 50.9, 65.0, 78.4, 113.0, 117.5, 122.5 (q, *J* = 275 Hz), 124.3, 124.9, 129.6, 130.2, 133.1 (q, *J* = 34 Hz), 133.3, 140.4, 148.6, 155.1 ppm; ¹⁹F NMR (282 MHz, δ, CDCl₃, 298 K): −63.0 ppm; IR (film): $\bar{\nu}$ = 3462, 3254, 3031, 2944, 2866, 1692, 1600, 1548, 1529, 1485, 1451, 1434, 1372, 1352, 1325, 1280, 1206, 1178, 1137, 904, 843, 830, 798, 737, 709, 683 cm^{−1}; HRMS (ESI): *m/z* calcd for C₂₄H₂₇F₆N₄O₃⁺: 533.1982 [M⁺], found: 533.1998.

General procedure for the asymmetric Michael reactions of 5

Degassed toluene (5 mL) was added to a mixture of the Schiff base 5 (0.1 mmol), catalyst **1c** (10 mol%), and Cs₂CO₃ (1.5 eq.) in a Schlenk tube. The stirring rate was set to 1000 rpm and the corresponding electrophile **15** (1.5 eq.) was added. After 24 h at 25 °C the reaction mixture was filtered through a plug of Na₂SO₄. The solvent was removed under reduced pressure. The crude products were purified by column chromatography (silica gel, heptanes : EtOAc = 20 : 1 to 2 : 1) giving the Michael addition products (**16**) in the reported yields.

R-(+)-16a. Obtained as a colourless oil in 85% yield (>95% conv.) and with an e.r. = 95 : 5 upon reacting Schiff base **5a** with acrylate **15a** in the presence of 10 mol% **1c** at 25 °C under the general procedure conditions. The analytical data are in full accordance with those reported in the literature.^{11,12} [α]_D²¹ (*c* = 0.70, CHCl₃) = 74.9°; ¹H NMR (300 MHz, δ , CDCl₃, 298 K): 1.44 (s, 9H), 2.16–2.27 (m, 2H), 2.33–2.41 (m, 2H), 3.59 (s, 3H), 3.93–3.99 (m, 1H), 7.14–7.21 (m, 2H), 7.28–7.47 (m, 6H), 7.60–7.68 (m, 2H) ppm; ¹³C NMR (75 MHz, δ , CDCl₃, 298 K): 28.2, 28.8, 30.5, 51.7, 64.9, 81.3, 127.9, 128.1, 128.6, 128.7, 128.9, 130.5, 136.6, 139.6, 170.8, 170.9, 173.7 ppm; IR (film): $\bar{\nu}$ = 2978, 2926, 1738, 1707, 1661, 1599, 1578, 1449, 1369, 1319, 1279, 1260, 1234, 1153, 943, 920, 849, 812 cm⁻¹; the enantioselectivity was determined using HPLC (Chiralcel AD-H, eluent: *n*-hexane : *i*-PrOH = 95 : 5, 0.5 mL min⁻¹, 10 °C, retention times: 10.9 min (major; *R*-enantiomer), 12.3 min (minor; *S*-enantiomer)); the absolute configuration was determined by comparison of the retention times and [α]_D value with literature data.¹² HRMS (ESI): *m/z* calcd for C₂₃H₂₇NO₄: 382.2013 [M + H]⁺; found: 382.2013.

General procedure for the cascade reactions of 5

In a round-bottom flask, a 2-cyanobenzaldehyde **6** (0.10 mmol) was added at room temperature to a stirred solution of glycine Schiff base **5** (1.1 eq., 0.11 mmol), K₂CO₃ (1 eq.), and catalyst **1d** (5% mol) in CH₂Cl₂ (3 mL). The mixture was stirred at r.t. for 24 h (1000 rpm). Afterwards, the mixture was purified directly using flash chromatography on silica gel with hexane : ethyl acetate = 8 : 2 to give the intermediate **7** as a mixture of diastereoisomers. The products (**7**) were dissolved in a cooled solution of 0.5 M HCl (1 mL) and THF (3 mL) (0 °C). The mixture was stirred at the same temperature for 2 h and then concentrated under vacuum. The resulting residue was treated with saturated NaHCO₃ (20 mL), extracted with CH₂Cl₂ (4 × 30 mL), and purified using flash chromatography (silica gel, hexanes : EtOAc = 2 : 1).

(-)-8a. Obtained as an amorphous solid in 83% yield (22 mg, 0.083 mmol) with a d.r. = 4 : 1 and e.r. = 95 : 5 for the major diastereoisomer. [α]_D²⁰ (*c* = 0.5, CHCl₃) = -2.5°. ¹H-NMR (300 MHz, δ , CDCl₃, 298 K): 1.40 (s, 9H), 1.61 (br s, 2H), 4.04 (d, *J* = 3.6 Hz, 1H), 5.77 (d, *J* = 3.5 Hz, 1H), 7.44 (d, *J* = 7.6 Hz, 1H), 7.58 (t, *J* = 7.42 Hz, 1H), 7.68 (t, *J* = 6.48 Hz, 1H), 7.92 (d, *J* = 7.6 Hz, 1H); ¹³C-NMR (100 MHz, δ , CDCl₃): 29.1, 58.7, 83.3, 83.9, 123.8, 126.9, 128.3, 130.8, 135.2, 147.5, 171.3, 171.5; MS (ESI): *m/z* = 264.1 (M + H)⁺; anal. calcd for C₁₄H₁₇NO₄: C, 63.87; H, 6.51; N, 5.32. Found: C, 63.97; H, 6.41; N, 5.37%. The enantioselectivity

was determined using HPLC (Chiralcel OD-H, eluent: *n*-hexane : *i*-PrOH = 90 : 10, 0.7 mL min⁻¹, 25 °C, retention times (major diastereomer): 25.3 min (minor), 34.3 min (major)).

Conclusions

A flexible and highly telescoped synthesis strategy for a structurally diverse library of chiral cyclohexanediamine-based urea-containing quaternary ammonium salts was developed. The catalysts were successfully employed in asymmetric Michael addition reactions and a new powerful aldol-initiated cascade reaction of glycine Schiff bases with cyanobenzaldehyde derivatives. In both cases the flexible catalyst strategy allowed us to systematically fine-tune the catalysts for these reactions, thus resulting in high enantioselectivities and good to excellent yields. Besides the high enantioselectivities, it was also shown that these catalysts are very promising for the control of *s-trans* Michael acceptors, thus providing a powerful catalyst platform for further challenging asymmetric transformations.

Acknowledgements

This work was supported by the Austrian Science Funds (FWF): Project No. P26387-N28. Financial support from the Federal State Government of Upper Austria (Research Fellowship to J. N.) is kindly acknowledged. We are grateful to Dr Markus Himmelsbach for his support with HRMS analysis. The NMR spectrometers used at JKU Linz were acquired in collaboration with the University of South Bohemia (CZ) with financial support from the European Union through the EFRE INTERREG IV ETC-AT-CZ program (project M00146, “RERI-uasb”).

Notes and references

- For selected reviews about asymmetric phase-transfer catalysis see: (a) K. Maruoka, *Asymmetric Phase Transfer Catalysis*, WILEY-VCH, Weinheim, 2008; (b) M. J. O'Donnell, in *Catalytic Asymmetric Syntheses*, ed. I. Ojima, WILEY-VCH, New York, 2nd edn, 2000, pp. 727–755; (c) K. Maruoka and T. Ooi, *Chem. Rev.*, 2003, **103**, 3013–3028; (d) M. J. O'Donnell, *Acc. Chem. Res.*, 2004, **37**, 506–517; (e) T. Ooi and K. Maruoka, *Angew. Chem., Int. Ed.*, 2007, **46**, 4222–4266; (f) S. Shirakawa and K. Maruoka, *Angew. Chem., Int. Ed.*, 2013, **52**, 4312–4348; (g) S.-S. Jew and H.-G. Park, *Chem. Commun.*, 2009, 7090–7103; (h) R. Herchl and M. Waser, *Tetrahedron*, 2014, **70**, 1935–1960.
- For a review about bifunctional chiral ammonium salt-based catalysts see: J. Novacek and M. Waser, *Eur. J. Org. Chem.*, 2013, 637–648.
- For pioneering reports using Cinchona alkaloid-based free OH-containing ammonium salt catalysts see: (a) R. Helder, J. C. Hummelen, R. W. P. M. Laane, J. S. Wiering and H. Wynberg, *Tetrahedron Lett.*, 1976, **17**, 1831–1834; (b) U. H. Dolling, P. Davis and E. J. J. Grabowski, *J. Am. Chem. Soc.*, 1984, **106**, 446–447; (c) M. J. O'Donnell, W. D. Bennett and S. Wu, *J. Am. Chem. Soc.*, 1989, **111**, 2353–2355.

- 4 For selected very recent examples describing the beneficial use of free OH-containing chiral ammonium salt catalysts see: (a) J. R. Wolstenhulme, A. Cavell, M. Gredicak, R. W. Driver and M. D. Smith, *Chem. Commun.*, 2014, **50**, 13585–13588; (b) B. Xiang, K. M. Belyk, R. A. Reamer and N. Yasuda, *Angew. Chem., Int. Ed.*, 2014, **53**, 8515–8518; (c) R. J. Armstrong and M. D. Smith, *Angew. Chem., Int. Ed.*, 2014, **53**, 12822–12826; (d) S. Shirakawa, L. Wang, R. He, S. Arimitsu and K. Maruoka, *Chem.–Asian J.*, 2014, **9**, 1586–1593; (e) S. Shirakawa, T. Tokuda, S. B. J. Kan and K. Maruoka, *Org. Chem. Front.*, 2015, **2**, 336–339.
- 5 For (thio)-urea containing Cinchona alkaloid-based ammonium salt catalysts see: (a) P. Bernal, R. Fernández and J. M. Lassaletta, *Chem.–Eur. J.*, 2010, **16**, 7714–7718; (b) K. M. Johnson, M. S. Rattley, F. Sladojevich, D. M. Barber, M. G. Nunez, A. M. Goldys and D. J. Dixon, *Org. Lett.*, 2012, **14**, 2492–2495; (c) M. Li, P. A. Woods and M. D. Smith, *Chem. Sci.*, 2013, **4**, 2907–2911; (d) B. Wang, Y. Liu, C. Sun, Z. Wei, J. Cao, D. Liang, Y. Lin and H. Duan, *Org. Lett.*, 2014, **16**, 6432–6435; (e) P. G. K. Clark, L. C. C. Vieira, C. Tallant, O. Fedorov, D. C. Singleton, C. M. Rogers, O. P. Monteiro, J. M. Bennett, R. Baronio, S. Müller, D. L. Daniels, J. Mendez, S. Knapp, P. E. Brennan and D. J. Dixon, *Angew. Chem., Int. Ed.*, 2015, **54**, 6217–6221.
- 6 For amino acid-based (thio)-urea containing ammonium salts see: (a) H.-Y. Wang, Z. Chai and G. Zhao, *Tetrahedron*, 2013, **69**, 5104–5111; (b) H.-Y. Wang, J.-X. Zhang, D.-D. Cao and G. Zhao, *ACS Catal.*, 2013, **3**, 2218–2221.
- 7 For *trans*-1,2-cyclohexanediamine-based (thio)-urea containing ammonium salts developed by our groups see: (a) J. Novacek and M. Waser, *Eur. J. Org. Chem.*, 2014, 802–809; (b) M. Perillo, A. Di Mola, R. Filosa, L. Palombi and A. Massa, *RSC Adv.*, 2014, **4**, 4239–4246.
- 8 T. Li, S. Zhou, J. Wang, J. L. Acena, V. A. Soloshonok and H. Liu, *Chem. Commun.*, 2015, **51**, 1624–1626.
- 9 (a) A. E. Sorochinsky, J. L. Acena, H. Moriwaki, T. Sato and V. A. Soloshonok, *Amino Acids*, 2013, **45**, 691–718; (b) A. E. Sorochinsky, J. L. Acena, H. Moriwaki, T. Sato and V. A. Soloshonok, *Amino Acids*, 2013, **45**, 1017–1033.
- 10 D. W. Lee, H.-J. Ha and W. K. Lee, *Synth. Commun.*, 2007, **37**, 737–742.
- 11 Use of TADDOL-based PTCs: G. N. Gururaja, T. Herchl, A. Pichler, K. Gratzer and M. Waser, *Molecules*, 2013, **18**, 4357–4372.
- 12 Use of cyclopropenimine-based chiral base catalysts:(a) J. Bandar and T. H. Lambert, *J. Am. Chem. Soc.*, 2012, **134**, 5552–5555; (b) J. Bandar, G. S. Sauer, W. D. Wulff, T. H. Lambert and M. J. Veticatt, *J. Am. Chem. Soc.*, 2014, **136**, 10700–10707; (c) J. Bandar, A. Barthelme, A. Y. Mazori and T. H. Lambert, *Chem. Sci.*, 2015, **6**, 1537–1547.
- 13 Use of oxo-pyrimidinium-based catalysts: A. E. Sheshenev, E. V. Boltukhina, A. J. P. White and K. K. Hii, *Angew. Chem., Int. Ed.*, 2013, **52**, 6988–6991.
- 14 In those cases where the conversion was at least 60–70% after 24 h, stirring for another day was usually sufficient to ensure >95% conversion of starting material 5.
- 15 R. Ceccarelli, S. Insogna and M. Bella, *Org. Biomol. Chem.*, 2006, **4**, 4281–4284.
- 16 For other reports on the synthesis of this compound see: (a) E. J. Corey, M. C. Noe and F. Xu, *Tetrahedron Lett.*, 1998, **39**, 5347–5350; (b) R. Chinchilla, P. Mazón, C. Nájera, F. J. Ortega and M. Yus, *ARKIVOC*, 2005, 222–232; (c) V. I. Maleev, M. North, V. A. Larionov, I. V. Fedyanin, T. F. Savel'yeva, M. A. Moscalenko, A. F. Smolyakov and Y. N. Belokon, *Adv. Synth. Catal.*, 2014, **356**, 1803–1810.
- 17 For a highly enantioselective but only moderately diastereoselective copper-catalyzed protocol for this reaction see: M. Strohmeier, K. Leach and M. A. Zajak, *Angew. Chem.*, 2011, **123**, 12543–12546.
- 18 For a comprehensive overview see: J. M. Seco, E. Quinoa and R. Riguera, *Chem. Rev.*, 2004, **104**, 17–117.
- 19 Detailed VCD studies to unambiguously prove this assignment will soon be carried out in collaboration with specialists.